

	<h2>Bariatric Surgery Guidelines</h2>	
<p>Guideline # 6174</p>	<p>Categories Clinical → Care Management CM, TCHP Guidelines, Utilization Management UM</p>	<p>This Guideline Applies To: Texas Children's Health Plan</p>
		<p>Document Owner Bhavana Babber</p>

GUIDELINE STATEMENT:

Texas Children's Health Plan (TCHP) performs authorization of all bariatric surgery.

DEFINITIONS:

Bariatric Surgery involves a variety of surgical weight loss procedures such as gastric bypass. The surgical procedure aims to modify the anatomy of the digestive system to help lose weight.

It is used to treat medical conditions that are caused or significantly worsened by obesity where comorbid conditions cannot be adequately treated by standard measures unless significant weight loss occurs.

PRIOR AUTHORIZATION GUIDELINES

1. Gastric procedures for weight loss (Bariatric surgery) are excluded from coverage for members covered by CHIP.
2. All requests for prior authorization for bariatric surgery are received via online submission, fax, phone or mail by the Utilization Management Department and processed during normal business hours.
3. The Utilization Management professional receiving the request evaluates the submitted information to determine if the documentation supports the bariatric surgery as an eligible service.
4. The physician requesting authorization for the surgery must provide documentation of **ALL** of the following:
 - 4.1. Documentation of unsuccessful medical treatment for obesity
 - 4.2. The individual's psychiatric profile is such that the candidate is able to understand, tolerate and comply with all phases of care and is committed to long-term follow-up requirements;

AND

- 4.3. The candidate's post-operative expectations have been addressed; **AND**
- 4.4. The individual has undergone a preoperative mental health assessment by a mental health professional and is felt to be an acceptable candidate if they have a history of psychiatric or psychological disorders, are under the care of a mental health professional or they are on psychotropic medications; **AND**
- 4.5. The individual has received a thorough explanation of the risks, benefits, and uncertainties of the procedure; **AND**
- 4.6. The candidate's treatment plan includes pre- and post-operative dietary evaluations and nutritional counseling; **AND**
- 4.7. The candidate's treatment plan includes counseling regarding exercise, psychological issues and the availability of supportive resources when needed.
- 4.8. Referral for bariatric surgery to the bariatric surgeon was completed by the practitioner who is treating the comorbid condition(s).
- 4.9. Identification of the facility where the services will be provided and attestation that it is a facility in Texas that is one of the following:
 - 4.9.1. Accredited by the Metabolic and Bariatric Surgery Accreditation and Quality Improvement Program (MBSAQIP).
 - 4.9.2. A children's hospital that has a bariatric surgery program and provides access to an experienced surgeon who employs a team that is capable of long-term follow-up of the metabolic and psychosocial needs of the member and family.
- 4.10. That **NONE** of the following additional contraindications exist:
 - 4.10.1 Endocrine cause for obesity, inflammatory bowel disease, chronic pancreatitis, cirrhosis, portal hypertension, or abnormalities of the gastrointestinal tract
 - 4.10.2 Chronic, long-term steroid treatment
 - 4.10.3 Pregnant, or plans to become pregnant within 18 months
 - 4.10.4 Noncompliance with medical treatment
 - 4.10.5 Significant psychological disorders that would be exacerbated or interfere with the long-term management of the member after the operation
 - 4.10.6 Active malignancy
- 5. Utilization Management professionals will reference the most recent available version of InterQual criteria to establish medical necessity for the bariatric surgery.
- 6. Repeat bariatric surgery may be considered medically necessary in either of the following circumstances:

6.1 To correct complications from bariatric surgery such as band malfunction, obstruction, or

stricture

- 6.2 To convert to a Roux-en-Y gastroenterostomy or to correct pouch failure in an otherwise compliant member when the initial bariatric surgery met medical necessity criteria
- 6.3 Conversion to a Roux-en-Y gastroenterostomy may be considered medically necessary for members who have not had adequate success (defined as a loss of more than 50 percent of excess body weight) two years following the primary bariatric surgery procedure, and the member has been compliant with a prescribed nutrition and exercise program following the procedure
7. Bariatric surgery is not medically necessary when the primary purpose of the surgery is any of the following:
 - 7.1. For weight loss for its own sake
 - 7.2. For cosmetic purposes
 - 7.3. For reasons of psychological dissatisfaction with personal body image
 - 7.4. For the member's or provider's convenience or preference
8. Requests that do not meet the criteria established by this guideline will be referred to a TCHP Medical Director/Physician Reviewer for review and the Denial Policy may be followed.
9. Preauthorization is based on medical necessity and not a guarantee of benefits or eligibility. Even if preauthorization is approved for treatment or a particular service, that authorization applies only to the medical necessity of treatment or service. All services are subject to benefit limitations and exclusions. Providers are subject to State and Federal Regulatory compliance and failure to comply may result in retrospective audit and potential financial recoupment.

REFERENCES:

Peer Reviewed Publications:

- Abbatini F, Capoccia D, Casella G, et al. Type 2 diabetes in obese patients with body mass index of 30-35 kg/m²: sleeve gastrectomy versus medical treatment. *Surg Obes Relat Dis.* 2012; 8(1):20-24.
- Adams TD, Davidson LE, Litwin SE, et al. Health benefits of gastric bypass surgery after 6 years. *JAMA.* 2012; 308(11):1122-1131.
- Adams TD, Gress RE, Smith SC, et al. Long-term mortality after gastric bypass surgery. *N Engl J Med.* 2007; 357(8):753-761.
- Alami RS, Morton JM, Schuster R, et al. Is there a benefit to preoperative weight loss in gastric bypass patients? A prospective randomized trial. *Surg Obes Relat Dis.* 2007; 3(2):141-145; discussion 145-146.
- Ali MR, Baucom-Pro S, Broderick-Villa GA, et al. Weight loss before gastric bypass: feasibility and effect on postoperative weight loss and weight loss maintenance. *Surg Obes Relat Dis.* 2007; 3(5):515-520.

- Angrisani L, Lorenzo M, Borrelli V. Laparoscopic adjustable gastric banding versus Roux-en-Y gastric bypass: 5-year results of a prospective randomized trial. *Surg Obes Relat Dis.* 2007; 3(2):127-132; discussion 132-133.
- Arias E, Martinez PR, Ka Ming V, et al. Mid-term follow-up after sleeve gastrectomy as a final approach for morbid obesity. *Obes Surg.* 2009; 19(5):544-548.
- Armstrong Sarah C., Bolling Christopher F., Michalsky Marc P., Reichard Kirk W. and section on obesity, section on surgery Pediatric Metabolic and Bariatric Surgery: Evidence, Barriers, and Best Practices Pediatrics December 2019, 144 (6) e20193223;
- Arterburn D, Powers JD, Toh S, et al. Comparative effectiveness of laparoscopic adjustable gastric banding vs laparoscopic gastric bypass. *JAMA Surg.* 2014; Oct 29 [Epub ahead of print].
- Ashrafian H, le Roux CW, Darzi A, Athanasiou T. Effects of bariatric surgery on cardiovascular function. *Circulation.* 2008; 118(20):2091-2102.
- Birkmeyer NJ, Dimick JB, Share D, et al. Hospital complication rates with bariatric surgery in Michigan. *JAMA.* 2010; 304(4):435-442.
- Bohdjalian A, Langer FB, Shakeri-Leidenmuhler S, et al. Sleeve gastrectomy as sole and definitive bariatric procedure: 5-year results for weight loss and ghrelin. *Obes Surg.* 2010; 20(5):535-540.
- Bowne WB, Julliard K, Castro AE, et al. Laparoscopic gastric bypass is superior to adjustable gastric band in super morbidly obese patients. *Arch Surg.* 2006; 141(7):683-689.
- Bray GA. The missing link – lose weight, live longer. *N Engl J Med.* 2007; 357(8):818-820.
- Brethauer SA, Hammel JP, Schauer PR. Systematic review of sleeve gastrectomy as staging and primary bariatric procedure. *Surg Obes Relat Dis.* 2009; 5(4):469-475.
- Brolin RE. Results of obesity surgery. *Gastrointestinal Clin N Am.* 1987; 16(2):317-335.
- Buchwald H, Avidor Y, Braunwald E, et al. Bariatric surgery: a systematic review and meta-analysis. *JAMA.* 2004; 292(14):1724-1737.
- Buchwald H, Estok R, Fahrenbach K, et al. Weight and type 2 diabetes after bariatric surgery: systematic review and meta-analysis. *Am J Med.* 2009; 122(3):248-256. e5.
- Bult MJ, van Dalen T, Muller AF. Surgical treatment of obesity. *Eur J Endocrin.* 2008; 158(2):135-145.
- Busetto L, Segato G, De Luca M, et al. Preoperative weight loss by intragastric balloon in super-obese patients treated with laparoscopic gastric banding: a case-control study. *Obes Surg.* 2004; 14(5):671-676.
- Carlsson LM, Peltonen M, Ahlin S, et al. Bariatric surgery and prevention of type 2 diabetes in Swedish obese subjects. *N Engl J Med.* 2012; 367(8):695-704.
- Catalano MF, Rudic G, Anderson AJ, Chua TY. Weight gain after bariatric surgery as a result of a large gastric stoma: endotherapy with sodium morrhuate may prevent the need for surgical revision. *Gastrointest Endosc.* 2007; 66(2):240-245.
- Chapman AE, Kiroff G, Game P, et al. Laparoscopic adjustable gastric banding in the treatment of obesity: a systematic literature review. *Surgery.* 2004; 135(3):326-351.
- Choi J, Digiorgi M, Milone L, et al. Outcomes of laparoscopic adjustable gastric banding in patients with low body mass index. *Surg Obes Relat Dis.* 2010; 6(4):367-371.
- Consensus Development Conference Panel. Gastrointestinal surgery for severe obesity. *Ann Int Med.* 1991; 115(12):956-961.
- DeMaria EJ. Bariatric surgery for morbid obesity. *N Engl J Med.* 2007; 356(21):2176-2183.
- DeMaria EJ, Winegar DA, Pate VW, et al. Early postoperative outcomes of metabolic surgery to treat diabetes from sites participating in the ASMBS bariatric surgery center of excellence program

as reported in the Bariatric Outcomes Longitudinal Database. *Ann Surg.* 2010; 252(3):559-566; discussion 566-567.

- Dietz WH, Robinson TN. Overweight children and adolescents. *N Engl J Med.* 2005; 352(20):2100-2109.
- Dixon AF, Dixon JB, O'Brien PE. Laparoscopic adjustable gastric banding induces prolonged satiety: a randomized blind crossover study. *J Clin Endocrinol Metab.* 2005; 90(2):813-819.
- Dolan K, Hatzifotis M, Newbury L, Fielding G. A comparison of laparoscopic adjustable gastric banding and biliopancreatic diversion in superobesity. *Obes Surg.* 2004; 14(2):165-169.
- Douketis JD, Feightner JW, Attia J, Feldman WF. Periodic health examination, 1999 update: 1. Detection, prevention and treatment of obesity. Canadian Task Force on Preventive Health Care. *CMAJ.* 1999; 160(4):513-525.
- Eissa MA, Gunner KB. Evaluation and management of obesity in children and adolescents. *J Pediatr Health Care.* 2004; 18(1):35-38.
- Encinosa WE, Bernard DM, Du D, Steiner CA. Recent improvements in bariatric surgery outcomes. *Med Care.* 2009; 47(5):531-535.
- Farrell TM, Haggerty SP, Overby DW, et al. Clinical application of laparoscopic bariatric surgery: an evidence-based review. *Surg Endosc.* 2009; 23(5):930-949.
- Fox SR, Fox KM, Srikanth MS, Rumbaut R. The Lap-Band system in a North American population. *Obes Surg.* 2003; 13(2):275-280.
- Frachetti KJ, Goldfine AB. Bariatric surgery for diabetes management. *Curr Opin Endocrinol Diabetes Obes.* 2009; 16(2):119-124.
- Fuks D, Verhaeghe P, Brehant O, et al. Results of laparoscopic sleeve gastrectomy: a prospective study in 135 patients with morbid obesity. *Surgery.* 2009; 145(1):106-113.
- Gandsas A, Li C, Tan M, et al. Initial outcomes following laparoscopic sleeve gastrectomy in 292 patients as a single-stage procedure for morbid obesity. *Bariatric Times.* 2010; 7(2):11-13.
- Gracia JA, Martínez M, Elia M, et al. Obesity surgery results depending on technique performed: long-term outcome. *Obes Surg.* 2009; 19(4):432-438.
- Garcia VF. Letters to the Editor: Adolescent bariatric surgery: treatment delayed may be treatment denied. *Pediatrics.* 2005; 115(3):822-823.
- Garcia VF, Langford L, Inge TH. Application of laparoscopy for bariatric surgery in adolescents. *Curr Opin Peds.* 2003; 15(3):248-255.
- Herron DM, Birkett DH, Thompson CC, et al. Gastric bypass pouch and stoma reduction using a transoral endoscopic anchor placement system: a feasibility study. *Surg Endosc.* 2008; 22(4):1093-1099.
- Hess DS, Hess DW, Oakley RS. The biliopancreatic diversion with the duodenal switch: results beyond 10 years. *Obes Surg.* 2005; 15(3):408-416.
- Himpens J, Dapri G, Cadière GB. A prospective randomized study between laparoscopic gastric banding and laparoscopic isolated sleeve gastrectomy: results after 1 and 3 years. *Obes Surg.* 2006; 16(11):1450-1456.
- Himpens J, Dobbelaire J, Peters G. Long-term results of laparoscopic sleeve gastrectomy for obesity. *Ann Surg.* 2010; 252(2):319-324.
- Hutter MM, Schirmer BD, Jones DB, et al. First report from the American College of Surgeons Bariatric Surgery Center Network: laparoscopic sleeve gastrectomy has morbidity and effectiveness positioned between the band and the bypass. *Ann Surg.* 2011; 254(3):410-422.
- Iannelli A, Dainese R, Piche T, et al. Laparoscopic sleeve gastrectomy for morbid obesity. *World J Gastroenterol.* 2008; 14(6):821-827.

- Inabnet WB, Quinn T, Gagner M, et al. Laparoscopic Roux-en-Y gastric bypass in patients with BMI < 50: a prospective randomized trial comparing short and long limb lengths. *Obes Surg.* 2005; 15(1):51-57.
- .
- Jamal MK, DeMaria EJ, Johnson JM, et al. Impact of major co-morbidities on mortality and complications after gastric bypass. *Surg Obes Relat Dis.* 2005; 1(6):511-516.
- Jan FC, Hong D, Pereira N, et al. Laparoscopic adjustable gastric banding versus laparoscopic gastric bypass for morbid obesity: a single institution comparison study of early results. *J Gastrointest Surg.* 2005; 9(1):30-41.
- Jossart GH, Cirangle PT. Laparoscopic sleeve gastrectomy: Mid-term weight loss results. *Bariatric Times.* 2009; S10-11.
- Karamanakos SN, Vagenas K, Kalfarentzos F, Alexandrikes TK. Weight loss, appetite suppression, and changes in fasting and postprandial ghrelin and peptide-YY levels after Roux-en-Y gastric bypass and sleeve gastrectomy: a prospective, double blind study. *Ann Surg.* 2008; 247(3):401-407.
- Lalor PF, Tucker ON, Szomstein S, Rosenthal RJ. Complications after laparoscopic sleeve gastrectomy. *Surg Obes Relat Dis.* 2008; 4(1):33-38.
- Langer FB, Bohdjalian A, Felberbauer FX, et al. Does gastric dilatation limit the success of sleeve gastrectomy as sole operation for morbid obesity? *Obes Surg.* 2006; 16(2):166-171.
- Lee CM, Cirangle PT, Jossart GH. Vertical gastrectomy for morbid obesity in 216 patients: report of two-year results. *Surg Endosc.* 2007; 21(10):1810-1816.
- Livhits M, Mercado C, Yermilov I, et al. Preoperative predictors of weight loss following bariatric surgery: systematic review. *Obes Surg.* 2012; 22(1):70-89.
- Lyass S, Cunneen SA, Hagiike M, et al. Device-related reoperations after laparoscopic adjustable gastric banding. *Am Surg.* 2005; 71(9):738-743.
- Marceau P, Hould FD, Simrad S, et al. Biliopancreatic diversion with duodenal switch. *World J Surg.* 1998; 22(9):947-954.
- Mason EE, Tang S, Renquist KE, et al. A decade of change in obesity surgery. National Bariatric Surgery Registry (NBSR) Contributors. *Obes Surg.* 1997; 7(13):189-197.
- McClean LD, Rhode BM, Nohr CW. Long- or short-limb gastric bypass? *J Gastrointest Surg.* 2001; 5(5):525-530.
- Meyer L, Rohr S, Becker J, et al. Retrospective study of laparoscopic adjustable silicone gastric banding for the treatment of morbid obesity: results and complications in 127 patients. *Diabetes Metab.* 2004; 30(1):53-60.
- Miller K, Hell E. Laparoscopic adjustable gastric banding: a prospective 4-year follow-up study. *Obes Surg.* 1999; 9(2):183-187.
- Mingrone G, Panunzi S, De Gaetano A, et al. Bariatric surgery versus conventional medical therapy for type 2 diabetes. *N Engl J Med.* 2012; 366(17):1577-1585.
- Mognol P, Chosidow D, Marmuse JP. Laparoscopic sleeve gastrectomy as an initial bariatric procedure for high risk patients: initial results in 10 patients. *Obes Surg.* 2005; 15(7):1030-1033.
- Mokdad AH, Ford ES, Bowman BA, et al. Prevalence of obesity, diabetes, and obesity-related health risk factors, 2001. *JAMA.* 2003; 289(1):76-79.
- Montgomery KF, Watkins BM, Ahroni JH, et al. Outpatient laparoscopic adjustable gastric banding in super-obese patients. *Obes Surg.* 2007; 17(6):711-716.
- Moon HS, Kim WW, Oh JH. Results of laparoscopic sleeve gastrectomy (LSG) at 1 year in morbidly obese Korean patients. *Obes Surg.* 2005; 15(10):1469-1475.

- Morino M, Toppino M, Bonnet G, del Genio G. Laparoscopic adjustable silicone gastric banding versus vertical banded gastroplasty in morbidly obese patients: a prospective randomized controlled clinical trial. *Ann Surg.* 2003; 238(6):835-842.
- Morton JM. Weight gain after bariatric surgery as a result of large gastric stoma: endotherapy with sodium morrhuate to induce stomal stenosis may prevent the need for surgical revision (editorial). *Gastrointest Endosc.* 2007; 66(2):246-247.
- Myers JA, Sarker S, Shayani V. Treatment of massive super-obesity with laparoscopic adjustable gastric banding. *Surg Obes Relat Dis.* 2006; 2(1):37-40.
- Neovius M, Narbro K, Keating C, et al. Health care use during 20 years following bariatric surgery. *JAMA.* 2012; 308(11):1132-1141.
- O'Brien PE, Dixon JB, Brown W, et al. The laparoscopic adjustable gastric band (Lap-Band): a prospective study of medium-term effects on weight, health and quality of life. *Obes Surg.* 2002; 12(5):652-660.
- O'Brien PE, Dixon JB, Laurie C, et al. Treatment of mild to moderate obesity with laparoscopic adjustable gastric banding or an intensive medical program. *Ann Intern Med.* 2006; 144(9):625-633.
- O'Brien PE, Dixon JB. Weight loss and early and late complications—the international experience. *Am J Surg.* 2002; 184(6 Suppl 2): S42-S45.
- O'Brien PE, Sawyer SM, Laurie C, et al. Laparoscopic adjustable gastric banding in severely obese adolescents: a randomized trial. *JAMA.* 2010; 303(6):519-526.
- Parikh MS, Ayoung-Chee P, Romanos E, et al. Comparison of rates of resolution of diabetes mellitus after gastric banding, gastric bypass, and biliopancreatic diversion. *J Am Coll Surg.* 2007; 205(5):631-635.
- Parikh M, Duncombe J, Fielding GA. Laparoscopic adjustable gastric banding for patients with body mass index of ≤ 35 kg/m². *Surg Obes Relat Dis.* 2006; 2(5):518-522.
- Parikh MS, Laker S, Weiner H, et al. Objective comparison of complications resulting from laparoscopic bariatric procedures. *J Am Coll Surg.* 2006; 202(2):252-261.
- Prachand VN, Davee RT, Alverdy JC. Duodenal switch provides superior weight loss in the superobese (BMI ≥ 50 kg/m²) compared with gastric bypass. *Ann Surg.* 2006; 244(4):611-619.
- Pratt JS, Lenders CM, Dionne EA, et al. Best practice updates for pediatric/adolescent weight loss surgery. *Obesity.* 2009; 17(5):901-910.
- Puzziferri N, Roshek TB 3rd, Mayo HG, et al. Long-term follow-up after bariatric surgery: a systematic review. *JAMA.* 2014; 312(9):934-942.
- Ramos A, Galvao Neto M, Galvao M, et al. Laparoscopic greater curvature plication: initial results of an alternative restrictive bariatric procedure. *Obes Surg.* 2010; 20(70):913-918.
- Regan JP, Inabnet WB, Gagner M, Pomp A. Early experience with two-stage laparoscopic Roux-en-Y gastric bypass as an alternative in the super-super obese patient. *Obes Surg.* 2003; 13(6):861-864.
- Romy S, Donadini A, Giusti V, Suter M. Roux-en-Y gastric bypass vs. gastric banding for morbid obesity: a case matched study of 442 patients. *Arch Surg.* 2012; 147(5):460-466.
- Rubenstein RB. Laparoscopic adjustable gastric banding at a U.S. center with up to 3-year follow-up. *Obes Surg.* 2002; 12(3):380-384.
- Rubin M, Yehoshua RT, Stein M, et al. Laparoscopic sleeve gastrectomy with minimal morbidity. Early results in 120 morbidly obese patients. *Obes Surg.* 2008; 18(12):1567-1570.

- Sanchez-Santos R, Masdevall C, Baltasar A, et al. Short- and mid-term outcomes of sleeve gastrectomy for morbid obesity: the experience of the Spanish National Registry. *Obes Surg.* 2009; 19(9):1203-1210.
- Schauer P, Chand B, Brethauer S. New applications for endoscopy: the emerging field of endoluminal and transgastric bariatric surgery. *Surg Endosc.* 2007; 21(3):347-356.
- Schauer PR, Kashyap SR, Wolski K, et al. Bariatric surgery versus intensive medical therapy in obese patients with diabetes. *N Engl J Med.* 2012; 366(17):1567-1576.
- Scopinaro N, Papadia F, Marinari G, et al. Long-term control of type 2 diabetes mellitus and the other major components of the metabolic syndrome after biliopancreatic diversion in patients with BMI < 35 kg/m². *Obes Surg.* 2007; 17(2):185-192.
- Silecchia G, Rizzello M, Casella G, et al. Two-stage laparoscopic biliopancreatic diversion with duodenal switch as treatment of high-risk super-obese patients: analysis of complications. *Surg Endosc.* 2009; 23(5):1032-1037.
- Sjöström L, Peltonen M, Jacobson P, et al. Bariatric surgery and long-term cardiovascular events. *JAMA.* 2012; 307(1):56-65.
- Sjöström L, Narbro K, Sjöström CD, et al. Effects of bariatric surgery on mortality in Swedish obese subjects. *N Engl J Med.* 2007; 357(8):741-752.
- Sjöström L, Peltonen M, Jacobson P, et al. Association of bariatric surgery with long-term remission of type 2 diabetes and with microvascular and macrovascular complications. *Clinical trials* No. NCT01479452. *JAMA.* 2014; 311(22):2297-2304.
- Spivak H, Beltran OR, Slavchev P, Wilson EB. Laparoscopic revision from LAP-BAND to gastric bypass. *Surg Endosc.* 2007; 21(8):1388-1392.
- Stanford A, Glascock JM, Eid GM, et al. Laparoscopic Roux-en-Y gastric bypass in morbidly obese adolescents. *J Pediatr Surg.* 2003; 38(3):430-433. Strain GW, Gagner M, Pomp A, et al. Comparison of weight loss and body composition changes with four surgical procedures. *Surg Obes Relat Dis.* 2009; 5(5):582-587.
- Sugerman JH, Sugerman EL, DeMaria EJ, et al. Bariatric surgery for severely obese adolescents. *J Gastrointest Surg.* 2003; 7(1):102-108.
- Sultan S, Parikh M, Youn H, et al. Early U.S. outcomes after laparoscopic adjustable gastric banding in patients with a body mass index less than 35 kg/m². *Surg Endosc.* 2009; 23(7):1569-1573.
- Thompson CC, Slattery J, Bundga ME, Lautz DB. Perioral endoscopic reduction of dilated gastrojejunal anastomoses after Roux-en-Y gastric bypass: a possible new option for patients with weight regain. *Surg Endosc.* 2006; 20(11):1744-1748.
- Treadwell JR, Sun F, Schoelles K. Systematic review and meta-analysis of bariatric surgery for pediatric obesity. *Ann Surg.* 2008; 248(5):763-776.
- Turner P. Slimming down for safer surgery. *Arch Surg.* 2009; 144(12):1155-1156.
- Weiner R, Bockhorn H, Rosenthal R, Wagner D. A prospective randomized trial of different laparoscopic gastric banding techniques for morbid obesity. *Surg Endosc.* 2001; 15(1):63-68.
- Zehetner J, Holzinger F, Triaca H, Klaiber CH. A 6-year experience with the Swedish adjustable gastric band: prospective long-term audit of laparoscopic gastric banding. *Surg Endosc.* 2005; 19(1):21-28.

Government Agency, Medical Society, and Other Publications:

GUIDELINE

- Uniform Managed Care Contract v2.28 Attachment B-2.1 – Medicaid and CHIP Managed Care Services RFP, CHIP Covered Services
- Texas Medicaid Provider Procedures Manual Medical and Nursing Specialists, Physicians, and Physician Assistants Handbook Accessed November, 2021
<https://www.tmhp.com/resources/provider-manuals/tmppm>
- American Society for Metabolic and Bariatric Surgery Pediatric Metabolic and Bariatric Surgery Guidelines published August 2018
<https://asmbs.org/resources/asmbs-pediatric-metabolic-and-bariatric-surgery-guidelines>

Status	Date	Action
Approved	1/13/2022	Clinical & Administrative Advisory Committee Reviewed and Approved for Implementation

Original Document Creation Date: 10/21/2016	This Version Creation Date: 01/04/21	Effective/Publication Date: 01/25/2022
---	--------------------------------------	--